



An evaluation of staining techniques for marking daily growth in scleractinian corals

Michael Holcomb^{a,b,*}, Anne L. Cohen^a, Daniel C. McCorkle^a

^a Dept. of Geology and Geophysics, Woods Hole Oceanographic Institution, Woods Hole, MA 02543, USA

^b Centre Scientifique de Monaco, Avenue Saint Martin, MC-98000 Monaco-Ville, Monaco

ARTICLE INFO

Article history:

Received 30 June 2012

Received in revised form 2 December 2012

Accepted 4 December 2012

Available online xxxx

Keywords:

Alizarin
Calcein
Calcification
Coral
Isotope
Stain

ABSTRACT

In situ skeletal markers have been widely used to quantify skeletal growth rates of scleractinian corals on sub-annual time-scales. Nevertheless, an evaluation of different techniques, both in terms of their efficacy and potential impacts on the growth process itself, has not been undertaken. Here the effects of exposure to four different dyes (alizarin, alizarin complexone, calcein, and oxytetracycline) and isotope spikes (Ba and Sr) on the growth rates of scleractinian corals are compared. Oxytetracycline increased coral growth. Alizarin, alizarin complexone, calcein, and Sr and Ba isotope spikes had no significant effect on coral growth, but polyp extension appeared reduced during exposure to alizarin and alizarin complexone. Calcein provided a more intense fluorescent mark than either alizarin or alizarin complexone. Isotope spikes were challenging to locate using isotope ratio analysis techniques. Thus, calcein appears best suited for marking short-term calcification increments in corals, while a combination of alizarin or alizarin complexone and calcein may be useful for dual labeling experiments as there is little overlap in their fluorescence spectra.

© 2012 Elsevier B.V. All rights reserved.

1. Introduction

Mounting concern about the impacts of climate change, ocean acidification and direct anthropogenic activities on coral reef ecosystems has spurred the need for accurate and precise quantification of the rates of skeletal growth of corals and other calcifying organisms, on diurnal through seasonal timescales, in both field and laboratory experiments. The most frequently used measure of skeletal growth rate in corals is based on annual banding, high and low density couplets that together represent one year. While extremely successful and widely used (e.g. Buddemeier, 1974; Cantin et al., 2010; Macintyre and Smith, 1974), this technique limits measures of growth to timescales of one year or longer, and cannot resolve sub-annual growth responses that are essential for understanding calcification responses to changes in light, temperature, nutrient availability, carbonate ion concentration, photosynthesis and catastrophic events such as storms.

A wide range of approaches have been developed to estimate coral growth on sub-annual timescales, including: alkalinity uptake (e.g. Jacques and Pilson, 1980; Smith, 1973), changes in buoyant weight (e.g. Davies, 1989), and radioisotope incorporation (e.g. Tambutte et al., 1995), direct physical measurement (e.g. Cruz-Pinon et al., 2003), time lapse photography (Barnes and Crossland, 1980), laser diffraction (Stromgren, 1976; Vago et al., 1997), and the use of various dye, elemental, and isotope spikes. Dye- and isotope-based approaches are commonly used in a range of calcifying organisms and offer many advantages

over other approaches. In particular, dye and isotope based marks are easily implemented in field settings, can be used in-situ, offer the ability to mark large numbers of organisms at the same time, and can be used on corals of vastly different sizes – from newly settled polyps to colonies meters across. Due to their ease of detection, dyes are commonly used to provide a time point marker within the coral skeleton as a means of estimating coral growth or identifying skeleton deposited within a particular time interval (Barnes, 1970, 1972; Cohen et al., 2004; Gladfelter, 1983; Marschal et al., 2004; Raz-Bahat et al., 2006; Tambutte et al., 2012; Venn et al., 2011). However, one commonly used dye, alizarin, has been shown to negatively affect the growth of corals (Dodge et al., 1984), thus alternative dyes are desirable. Here, four dyes (alizarin, alizarin complexone, calcein, and oxytetracycline) and stable isotope spikes (commonly used to mark fish otoliths (Thorrold et al., 2002, 2006) and bones (Sun et al., 1992)) were used to mark the skeletons of corals to assess if they were effective in marking coral skeletons and if exposure to the dye impacted coral growth. Absorption spectra for each dye in seawater are presented to assist in choosing dyes which do not absorb light in regions of the spectrum which may be of experimental interest. Emission spectra for each dye incorporated into the skeleton are also presented to aid in choosing appropriate filter sets for imaging the dyes with fluorescence microscopy.

2. Methods

2.1. Dyes

One of four dyes was used in each dye incubation to mark the skeleton: alizarin red S (sodium salt – Alfa Aesar 42040 lot E22R017 –

* Corresponding author at: University of Western Australia, School of Earth and Environment, 35 Stirling Highway, 6009 Crawley, WA, Australia. Tel.: +377 11 208 475 4249. E-mail address: mholcomb3051@gmail.com (M. Holcomb).

referred to as alizarin throughout this manuscript), alizarin complexone (Alfa Aesar A16699 lot E8180A), calcein (Alfa Aesar L10255 lot USLF006789 – this particular lot was soluble in distilled water, suggesting that it was in the form of salt), and oxytetracycline HCl (USB 23659 lot 113648). In addition, isotope spikes (Ba 135 or Sr 86, purchased as carbonate salts from Oak Ridge National Lab) were used as markers in some incubations.

All dyes and isotope spikes were added as aliquots of stock solutions. Spikes were mixed with sufficient HCl to dissolve the carbonate salt and made up in distilled water to make stock solutions of which 50–125 μ l was used per liter of seawater. For Sr isotope spikes, 50 μ l of a given stock solution added to a liter of seawater doubled the concentration of that particular Sr isotope. For Ba isotope spikes, 50 μ l of stock solution almost doubled the total Ba concentration.

2.2. Coral maintenance

Colonies of the temperate scleractinian coral *Astrangia poculata* were collected and processed as described by Holcomb et al. (2010), except that in addition to colonies, newly settled polyps and their associated substratum were also attached to slides. All slides with corals were suspended vertically in a flow-through aquarium receiving filtered (20 μ m) Vineyard Sound seawater. Incoming seawater was heated in the winter, thus corals experienced a temperature range of 14–30 °C; temperatures at the time of experiments are as specified. Aquaria were aerated to maintain water circulation. Corals were maintained under aquarium conditions for at least one month prior to use in experiments. A mixture of brown and white colonies (zooxanthellate and azooxanthellate colonies) each ~2–5 cm in diameter was used for all treatments. Corals were fed regularly with newly hatched and frozen brine shrimp.

For marking experiments, corals were placed in pre-washed (with fresh and seawater) 1 l PET food service containers with lids (SOLO) containing ~800 ml of water from the source aquarium. Airstones were added to each container and each container bubbled continuously. Containers were held within a water bath with a temperature similar to that of the source aquarium.

2.3. Dyeing corals

In dye experiments with *A. poculata* (March–Oct. 2009), growth rates were estimated via alkalinity depletion measurements the day before (pre-treatment), the day of (treatment), and the day after (post-treatment) dye exposure. All alkalinity incubations were ~24 h in duration, covering a full light–dark cycle. The temperature range was 25–26 °C. For each treatment 4–7 corals were used, each in an individual incubation container. At the same time as dye treatments, additional corals not exposed to dye were also measured to control for day-to-day variations in growth. Incubations were carried out in 1 l PET food service containers: ~800 ml (actual amount weighed to 0.01 g) of water from the source tank was added to each container, and a coral was added. Containers with no coral added were used to estimate background changes in alkalinity. Irradiance (PAR – measured with a diving-PAM underwater quantum sensor (WALZ)) ranged from 10 to 40 μ mol photons/m²/s with a 12 h light dark cycle (white colonies were incubated under the lower end of the range of light levels, and brown colonies under the higher light levels – a similar light gradient was present in the source tank due to different corals being at different distances from the light bulbs or being closer to the ends of the light bulbs which produce less light than the center). Light was provided by two T5-HO bulbs (10,000 K, 54 W).

Alkalinity samples were taken from each container ~1 h after the corals had been added and again at the end of the incubation. Waiting 1 h after the addition of the coral to take the first sample was intended to allow the coral to recover from any handling stress and thus avoid capturing any temporary changes in calcification. Salinity

(Hach conductivity probe – read to 0.1, accurate to ~1) and pH (NBS scale, Thermo-Orion ROSS 8165BNWP electrode, read to 0.1 mV) were measured at the end of each incubation for every container, as well as at the start of incubations for a subset of the containers. Aragonite deposition was assumed to be the only process affecting alkalinity, with 2 mol alkalinity consumed per mol of CaCO₃ deposited. This may under-estimate calcification as any ammonia released by the coral will increase the alkalinity of the solution (e.g. Jacques and Pilson, 1980). Alkalinity depletion rates were corrected for evaporation (based on the change in container mass), and for background rates measured in containers containing no slides. Background alkalinity consumption rates were invariably low, with the highest rates being <10% of coral rates.

Final dye concentrations were as follows: 2.7–3.2 mg/kg alizarin (added as ~0.2 ml of stock solution/l, pH was not adjusted, but pH declined <0.01 upon dye addition), 8.6–8.8 mg/kg alizarin complexone (added as ~1 ml of stock solution/l with sufficient NaOH to dissolve, pH declined ~0.03 upon dye addition), 9.5–10 mg/kg calcein (added as ~0.8 ml of stock solution/l, pH of the stock solution was not adjusted, thus pH declined ~0.03 upon dye addition), 24–26 mg/kg oxytetracycline (added as ~0.3 ml/l of stock suspension, pH was adjusted with NaOH, no measureable pH change upon addition).

2.4. Isotope spikes

Marking corals with isotope spikes was carried out as a part of long term growth experiments (see Holcomb et al., 2010, 2012); data from control corals included in those experiments are presented here. Isotope experiments were carried out using two different isotopes, ⁸⁶Sr and ¹³⁵Ba, with 6–16 corals for each treatment. Marking with ⁸⁶Sr was carried out in much the same manner as that used for dye experiments, with 60 μ l of an ⁸⁶Sr solution added to ~800 ml seawater and corals were incubated for two days. Growth was estimated from changes in buoyant weight (per Holcomb et al., 2010) for the 5 months prior to and the month following the isotope spike. Corals were held at one of two temperatures – ~19 or ~26 °C throughout that six month period.

Spikes with ¹³⁵Ba were carried out in a flow-through aquarium system as used by Holcomb et al. (2012). Each reservoir used to supply water to individual aquaria was spiked with 81 μ l of ¹³⁵Ba solution/l of seawater. Individual aquaria received spiked seawater for a period of two days: unspiked seawater was then added to the reservoir, diluting the spike to ~80%, and each subsequent day the remaining spike was diluted by an additional ~60%. Buoyant weights were measured for the two months prior and one month following isotope exposure using a Sartorius G803S balance, and the aquaria were held at either 16 or 24 °C throughout this period.

2.5. Alkalinity

Alkalinity samples were taken in pre-cleaned glass or plastic scintillation vials with screw top lids and foamed polyethylene liners (Wheaton). Samples were stored refrigerated for no more than 1 month prior to measurement. Alkalinities were measured via titration with 0.01 N HCl containing 40.7 g NaCl/l using a Metrohm Titrando 808 Dosimat and a 730 Sample Changer controlled by Tiamo software to perform automated normalized Gran titrations of 1 ml samples. Duplicate samples were run and additional replicates were run if measured values differed by more than 4 μ mol/kg. Certified seawater reference material supplied by the lab of Andrew Dickson (Scripps Institution of Oceanography) was run each time samples were run.

2.6. Spectra

2.6.1. Absorbance spectra

To estimate the potential effect of the presence of the dye on the light spectra received by the coral, the absorbance spectra of each

dye in seawater was measured using an Ocean Optics USB4000 spectrophotometer configured for measuring the visible spectrum with a blue filtered (SCHOTT glass BG-34 filter) tungsten light source (LS-1) and a 1 cm cuvette. In addition to the dyes used for the coral experiments, the absorption spectra were also measured for other lots of calcein (from Alfa Aesar, Invitrogen, and Sigma) and oxytetracycline (Acros). Different lots of calcein were found to be highly variable in their appearance and solubility – some being readily soluble in distilled water, while others requiring addition of NaOH to dissolve. Even when purchased as sodium salt, not all lots were soluble in distilled water. Thus when preparing calcein stock solutions the manner of preparation will depend greatly on the lot of calcein used. It may be possible to make up the solution directly in water, or it may require the use of a base, such as NaOH or NaHCO_3 (per Wilson et al., 1987) to solubilize the calcein; it is advisable to check the pH of the calcein stock solutions and adjust as needed before use.

2.6.2. Emission spectra

A Leica TCS SP5 confocal microscope at the Centre Scientifique de Monaco was used to measure the emission spectra of each dye incorporated into the coral skeleton. Polished sections of dyed coral skeletons were prepared using standard methods with water or mineral oil used to suspend polishing compounds (e.g. Holcomb et al., 2009). Fluorescence was excited using one of three lasers: a 543 nm He/Ne laser, a 488 nm Ar laser, or a 405 nm diode laser. The resulting fluorescence spectra were captured with the confocal microscope.

2.7. Statistics

All data for dye comparisons were expressed as relative growth rates – the ratio of the post treatment growth rate to the pretreatment growth rate – for statistical and plotting purposes (see Holcomb et al., 2012 for a discussion of this normalization approach).

$$\text{relative growth} = \frac{\frac{\Delta \text{massposttreatment}}{\Delta \text{timeposttreatment}}}{\frac{\Delta \text{masspretreatment}}{\Delta \text{timepretreatment}}}$$

Rates were further corrected for day to day changes in calcification by dividing by the average relative growth rate of untreated corals run at the same time. Similar patterns were observed for both brown and white colonies, thus data were pooled.

Differences among treatments were detected using a Kruskal–Wallis test (Systat 9), and if warranted, nonparametric multiple comparisons were performed to compare treatments to controls as per Zar (1984). For isotope treatments at different temperatures (Fig. 2), growth rates were normalized to starting mass; a sign test (Zar, 1984) was used to test whether the post isotope treatment growth rate differed from the pre-treatment rate at each temperature.

3. Results

All dyes used gave detectable marks in the skeleton and all corals survived the exposure. Growth rates of *A. poculata* were reduced during exposure to all dye treatments except oxytetracycline; rates during exposure were 84%, 77% and 70% of the initial rates for alizarin, alizarin complexone and calcein respectively (Fig. 1). Growth rates during oxytetracycline treatment were higher (114%) than the initial rates. Following dye exposure, growth rates returned to near pre-treatment rates – 99%, 125%, and 115% of the initial rates for alizarin, alizarin complexone and calcein respectively. Growth rates following oxytetracycline exposure, however, were significantly ($p < 0.01$) higher (168%) than the pretreatment rates. Exposure to isotope spikes had no measurable effect on coral growth (Fig. 2).

Absorption spectra (Fig. 3) measured in seawater showed peak absorbances for oxytetracycline at ~380 nm, calcein at ~486 nm,

alizarin at ~540 nm, and alizarin complexone at ~550 nm. Different lots of calcein and oxytetracycline had similar spectra, despite differences in solubility and appearance.

The emission spectra (Fig. 4) of dyes incorporated into coral skeletons showed peaks at ~550 nm for oxytetracycline when excited at 405 nm, ~520 nm for calcein when excited at 488 nm, ~610 nm for alizarin and ~630 nm for alizarin complexone when excited at 543 nm.

4. Discussion

All dyes employed in this study proved effective in marking coral skeletons. However, consistent with previous reports of alizarin negatively affecting growth (Dodge et al., 1984), growth rates tended to be slightly lower following exposure to alizarin (Fig. 1), and, though not quantified, polyps appeared less expanded during exposure to alizarin. Oxytetracycline significantly increased coral growth rates (Fig. 1C). Though it is not known why oxytetracycline increases growth rates, corals, such as *A. poculata*, are often host to a wide range of endolithic organisms which erode the skeleton from within (e.g. Tribollet et al., 2009). Oxytetracycline may negatively impact some of the boring organisms and thus could decrease dissolution rates, thereby increasing net calcification. Calcein, alizarin complexone and isotope spikes had no measurable effect (Figs. 1, 2), and for calcein and isotope spikes, there was no apparent difference in polyp behavior during exposure.

4.1. Isotopes

A few studies have used stable isotope markers in calcium carbonates (Holcomb et al., 2009; Houlbreque et al., 2008; Ries et al., 2010; Thorrold et al., 2006), but this approach is not common due both to the expense of the spikes and to the difficulty of detecting the spikes once in the skeleton, which generally requires specialized instruments such as secondary ion or laser ablation mass spectrometers. In principle, however isotope based techniques for marking coral skeletons offer several advantages. The use of isotopes with low natural abundances can allow a relatively small change in the concentration of a given element to yield a large change in the isotopic composition. Isotope spikes had no measurable effect on coral growth (Fig. 2), and since the elements used occur naturally in the skeleton and surrounding seawater, a small change in their concentration would not be expected to affect growth. Radio-isotopes have been used (Bonham, 1965; Knutson et al., 1972; Marshall and Wright, 1998) and the detection means are relatively accessible. However, the risks associated with artificially spiking corals with radioisotopes, and regulatory concerns, may make their use undesirable. The use of stable isotopes avoids these problems, and, since there are multiple stable isotopes available for many of the elements found in coral skeletons, it is possible to introduce distinct isotopic markers at different time points. Our results suggest that even with specialized instrumentation, dyes are still more convenient than stable isotopes for mapping calcification at high spatial resolution over large areas of the skeleton (see supplementary figures). Depending on which salt is purchased (e.g. carbonate or chloride), isotope spikes may require dissolution in acid and pH adjustment prior to use.

4.2. Dyes

Dye based methods have the potential to change growth due to the introduction of a foreign substance (e.g. Ibsen and Birkedal, 2010), and due to their inherent absorption of light, they will change the light spectrum (Fig. 3) received by the coral during the incubation, potentially affecting growth (e.g. Kinzie et al., 1984). Dyes have the potential to have long-term effects on the reflected light spectrum and thus may alter an important source of light for photosynthesis (e.g. Enriquez et al., 2005; Falkowski et al., 1990; Kuhl et al., 1995).




Fig. 1. Relative growth rates of corals versus treatment based on changes in alkalinity depletion measured during dye exposure (A), and following dye exposure (B). Bars represent average values for 11 untreated corals, 7 for alizarin, 4 each for alizarin complexone and calcein, and 5 for oxytetracycline. Treatments are as follows: None = untreated control, Aliz = alizarin, AC = alizarin complexone, Cal = calcein, and OTC = oxytetracycline. Bars are means, and error bars are standard deviation. The treatment significantly ($p < 0.01$) different from the control (OTC in B) is indicated by “**”.

Despite their limitations, dyes also offer many advantages. A range of dyes are available, so it is possible to introduce multiple time markers, and the dyes are all visualized with widely available techniques – visible light (alizarin, alizarin complexone) or fluorescence (alizarin, alizarin complexone, calcein, oxytetracycline) microscopy (see Supplemental Figs. 1, 2), and dye methods require relatively little specimen preparation.

4.3. Alizarin

Alizarin has a long history of use for marking coral skeletons and has been used on a wide range of species (e.g. Barnes, 1970, 1972; Cohen et al., 2004; Trotter et al., 2011). Alizarin is readily available as a sodium salt making solution preparation simple and reducing the need for pH adjustment. We have used various lots of alizarin and the dye appears fairly consistent lot to lot. The pink color of alizarin incorporated into the skeleton is readily distinguished from the unstained skeleton under visible light. Alizarin can be detected with fluorescence microscopy too, and its absorption and emission spectra are sufficiently different from those of calcein and oxytetracycline that it can potentially be used for dual labeling experiments. The pink-purple color of water containing alizarin allows its presence to be readily verified.




Fig. 2. Growth rates normalized to skeletal dry weight for *A. poculata* specimens treated with isotope spikes at different temperatures. At 26 °C and 19 °C, corals were treated with ⁸⁶Sr, 16 corals were used at 26 °C, and 6 at 19 °C. At 24 °C and 16 °C, corals were treated with ¹³⁵Ba, 10 corals were used at 24 °C, and 8 at 16 °C. Black bars are pre-treatment rates, gray bars are post treatment rates, values are means, and error bars are standard deviation. Post-treatment growth rates did not significantly ($p < 0.01$) differ from pre-treatment rates.

Unfortunately, alizarin can also negatively impact the growth of corals (e.g. Dodge et al., 1984). Our own observations suggest that 12 h exposures to 10 mg of alizarin/l of seawater can kill some species of corals and reduce polyp extension in other species yet have still no effect on others (we have since used <5 mg of alizarin/l of seawater and at this dye concentration have not observed any mortalities for a range of species). Alizarin will precipitate out of seawater if present at high concentrations (Barnes, 1972), thus when adding concentrated stock solutions the seawater must be vigorously mixed to prevent the formation of precipitates.

4.4. Alizarin complexone

Alizarin complexone is similar in many respects to alizarin, with the advantage that it has not been found to reduce the growth of any coral species, although our results suggest that it may reduce polyp extension in *A. poculata*. Unfortunately, alizarin complexone is not as easy to distinguish from unmarked skeleton using transmitted light microscopy – the purple coloration of alizarin complexone provides nowhere near the contrast of alizarin. However, it can be seen readily with fluorescence microscopy (see Supplementary materials), with a spectrum similar to that of alizarin. We have only used a single lot of alizarin complexone; for that lot a base must be used to get the alizarin complexone into solution, and pH adjustment is advisable.




Fig. 3. Absorption spectra for each dye in seawater. Alizarin (Aliz) 6.6 mg/kg, alizarin complexone (AC) 4.9 mg/kg, calcein (Cal) 8 mg/kg, and oxytetracycline (OTC) 26 mg/kg.




Fig. 4. Fluorescence emission spectra from coral skeletons (*Porites* and *Goniastrea*) containing different dyes, as well as background spectra taken on adjacent unstained regions of the skeleton, with fluorescence excited using different wavelengths. A. 405 nm excitation. B. 488 nm excitation. C. 543 nm excitation (some spectra also include 488 nm excitation). Note, the portion of the spectra within 10 nm of the excitation wavelength(s) has been deleted for clarity. Spectra are shown for OTC = oxytetracycline, BKG = background, taken on an unstained region of the coral, Cal = calcein, AC = alizarin complexone, CalAliz = calcein and alizarin staining the same region, Aliz = alizarin, Cal Aliz AC = calcein, alizarin, and alizarin complexone staining the same region.

4.5. Calcein

Calcein does not appear to affect coral growth and has been recommended over alizarin and Sr marking for some shellfish species (Herrmann et al., 2009; Riascos et al., 2007), though it may affect the growth of some organisms (Thebault et al., 2006). The effect of calcein on the incorporation of Sr and Mg into calcite has been investigated, and it has been found not to significantly affect the incorporation of these elements (Dissard et al., 2009). The brilliant yellow-green color of calcein in seawater makes it easy to detect when it is present. The formation of precipitates was not observed to be a problem, making it convenient to inject a concentrated stock solution into bags surrounding corals for field marking. Fluorescence microscopy is

required to detect calcein in the skeleton as the yellow-orange color of aragonite containing calcein is difficult to see, while the intense fluorescence of calcein is easily detected. Unfortunately there appears to be considerable variability between suppliers, and for certain suppliers, lot-lot variability in the composition of the material sold as calcein. So depending upon the lot, a base may be needed to dissolve the calcein or it may be readily soluble in distilled water, and pH adjustment may or may not be needed.

4.6. Oxytetracycline

Oxytetracycline suffers from relatively low solubility at seawater pH making the preparation of concentrated stock solutions more difficult, and if used without pH adjustment, it will reduce pH. The increase in growth observed following oxytetracycline exposure suggests that it affects the coral holobiont, and should be used with caution.

Of the dyes used, calcein appears to be the most satisfactory for marking coral skeletons. Calcein had no detectable effect (negative or positive) on coral growth and no obvious effect on tissue expansion, it is readily available, and relatively soluble (in alkaline solutions). The brilliant yellow-green color of the water following its addition makes it easy to verify that it is present in field settings, and its strong fluorescence signal allows it to be easily detected. Additionally, calcein has the potential to be used with alizarin or alizarin complexone to carry out dual marking studies.

Acknowledgments

Many thanks to A. Hoggett and L. Vail for making the work at Lizard Island Research Station possible, to M. Pearce, L. Pearce, M. Gagliano, H. Salomonsen, A. Heenan, D. Bayley, and others at LIRS, to M. Scott, K. Goudkamp and the Great Barrier Marine Park Authority for permitting assistance, and to K. Casciotti, G. Gaetani, L. Kerr, N. Shimizu, S. Tambutté, N. Techer, and A. Venn for use of equipment and assistance, to A. Bodendorf, S. Gallager, F. Keller, H. Keys, E. Roosen, A. Tarrant, K. Thompson and A. York for assistance, and to M. Brosnahan, L. Hmelo, C. Mingione, B. Pedler, C. Saenger, N. Trowbridge, K. Whalen, and T. Rioux for diving support. Funding for this work was provided by a Lizard Island Doctoral Fellowship, the Ocean Life Institute, NSF OCE-1041106, and an International Society for Reef Studies/Ocean Conservancy Fellowship. This material is based upon work supported under a National Science Foundation Graduate Research Fellowship and a National Science Foundation International Post-Doctoral Fellowship. Research conducted by the Centre Scientifique de Monaco is supported by the Principality of Monaco. [SS]

Appendix A. Supplementary data

Supplementary data to this article can be found online at <http://dx.doi.org/10.1016/j.jembe.2012.12.003>.

References

- Barnes, D.J., 1970. Coral skeletons: an explanation of their growth and structure. *Science* 170, 1305–1308.
- Barnes, D.J., 1972. The structure and formation of growth-ridges in scleractinian coral skeletons. *Proc. R. Soc. Lond. B* 182, 331–350.
- Barnes, D.J., Crossland, C.J., 1980. Diurnal and seasonal variations in the growth of a staghorn coral measured by time-lapse photography. *Limnol. Oceanogr.* 25, 1113–1117.
- Bonham, K., 1965. Growth rate of giant clam *Tridacna gigas* at Bikini Atoll as revealed by radioautography. *Science* 149, 300–302.
- Buddemeier, R.W., 1974. Environmental Controls over Annual and Lunar Monthly Cycles in Hermatypic Coral Calcification: *Proc. 2nd Int. Coral Reef Sym.*, 2, pp. 259–267.
- Cantin, N.E., Cohen, A.L., Karnauskas, K.B., Tarrant, A.M., McCorkle, D.C., 2010. Ocean warming slows coral growth in the central Red Sea. *Science* 329, 322–325.
- Cohen, A.L., Smith, S.R., McCartney, M.S., van Etten, J., 2004. How brain coral record climate: an integration of skeletal structure, growth and chemistry of *Diporia labyrinthiformis* from Bermuda. *Mar. Ecol. Prog. Ser.* 271, 147–158.

- Cruz-Pinon, G., Carricart-Ganivet, J.P., Espinoza-Avalos, J., 2003. Monthly skeletal extension rates of the hermatypic corals *Montastraea annularis* and *Montastraea faveolata*: biological and environmental controls. *Mar. Biol.* 143, 491–500.
- Davies, P.S., 1989. Short-term growth measurements of corals using an accurate buoyant weighing technique. *Mar. Biol.* 101, 389–395.
- Dissard, D., Nehrke, G., Reichart, G.J., Nouet, J., Bijma, J., 2009. Effect of the fluorescent indicator calcein on Mg and Sr incorporation into foraminiferal calcite. *Geochim. Geophys. Geosyst.* 10, Q11001. <http://dx.doi.org/10.1029/2009GC002417>.
- Dodge, R.E., Wyers, S.C., Frith, H.R., Knap, A.H., Smith, S.R., Cook, C.B., Sleeter, T.D., 1984. Coral calcification rates by the buoyant weight technique: effects of alizarin staining. *J. Exp. Mar. Biol. Ecol.* 75, 217–232.
- Enriquez, S., Mendez, E.R., Iglesias-Prieto, R., 2005. Multiple scattering on coral skeletons enhances light absorption by symbiotic algae. *Limnol. Oceanogr.* 50, 1025–1032.
- Falkowski, P.G., Jokiel, P.L., Kinzie, R.A.I., 1990. Irradiance and corals. *Coral Reefs ed Z Dubinsky*, pp. 89–107.
- Gladfelter, E.H., 1983. Skeletal development in *Acropora cervicornis* II. Diel patterns of calcium carbonate accretion. *Coral Reefs* 2, 91–100.
- Herrmann, M., Lepore, M.L., Laudien, J., Arntz, W.E., Penchaszadeh, P.E., 2009. Growth estimations of the Argentinean wedge clam *Donax hanleyanus*: a comparison between length–frequency distribution and size-increment analysis. *J. Exp. Mar. Biol. Ecol.* 379, 8–15.
- Holcomb, M., Cohen, A.L., Gabitov, R.I., Hutter, J.L., 2009. Compositional and morphological features of aragonite precipitated experimentally from seawater and biogenically by corals. *Geochim. Cosmochim. Acta* 73, 4166–4179.
- Holcomb, M., McCorkle, D.C., Cohen, A.L., 2010. Long-term effects of nutrient and CO₂ enrichment on the temperate coral *Astrangia poculata* (Ellis and Solander, 1786). *J. Exp. Mar. Biol. Ecol.* 386, 27–33.
- Holcomb, M., Cohen, A.L., McCorkle, D.C., 2012. An investigation of the calcification response of the scleractinian coral *Astrangia poculata* to elevated pCO₂ and the effects of nutrients, zooxanthellae and gender. *Biogeosciences* 9, 29–39. <http://dx.doi.org/10.5194/bg-9-29-2012>.
- Houlbreque, F., Meibom, A., Cuif, J.-P., Stolarski, J., Marrocchi, Y., Ferrier-Pages, C., Domart-Coulon, I., Dunbar, R.B., 2008. Strontium-86 labeling experiments show spatially heterogeneous skeletal formation in the scleractinian coral *Porites porites*. *Geophys. Res. Lett.* 36, L04604. <http://dx.doi.org/10.1029/2008GL036782>.
- Ibsen, C.J.S., Birkedal, H., 2010. Modification of bone-like apatite nanoparticle size and growth kinetics by alizarin red S. *Nanoscale*. <http://dx.doi.org/10.1039/c0nr00488j>.
- Jacques, T.G., Pilson, M.E.Q., 1980. Experimental ecology of the temperate scleractinian coral *Astrangia danae* I. Partition of respiration, photosynthesis and calcification between host and symbionts. *Mar. Biol.* 60, 167–178.
- Kinzie, R.A., Jokiel, P.L., York, R., 1984. Effects of light of altered spectral composition on coral zooxanthellae associations and on zooxanthellae in vitro. *Mar. Biol.* 78, 239–248.
- Knutson, D.W., Buddemeier, R.W., Smith, S.V., 1972. Coral chronometers: seasonal growth bands in reef corals. *Science* 177, 270–272.
- Kuhl, M., Cohen, Y., Dalsgaard, T., Jorgensen, B.B., Revsbech, N.P., 1995. Microenvironment and photosynthesis of zooxanthellae in scleractinian corals studied with microsensors for O₂, pH and light. *Mar. Ecol. Prog. Ser.* 117, 159–172.
- Macintyre, I.G., Smith, S.V., 1974. X-radiographic Studies of Skeletal Development in Coral Colonies: *Proc. 2nd Int. Coral Reef Sym.*, 2, pp. 277–287.
- Marschal, C., Garrabou, J., Harmelin, J.G., Pichon, M., 2004. A new method for measuring growth and age in the precious red coral *Corallium rubrum* (L.). *Coral Reefs* 23, 423–432.
- Marshall, A.T., Wright, A., 1998. Coral calcification: autoradiography of a scleractinian coral *Galaxea fascicularis* after incubation in ⁴⁵Ca and ¹⁴C. *Coral Reefs* 17, 37–47.
- Raz-Bahat, M., Erez, J., Rinkevich, B., 2006. In vivo light-microscopic documentation for primary calcification processes in the hermatypic coral *Stylophora pistillata*. *Cell Tissue Res.* 325, 361–368.
- Riascos, J.M., Guzman, N., Laudien, J., Heilmayer, O., Oliva, M., 2007. Suitability of three stains to mark shells of *Concholepas concholepas* (Gastropoda) and *Mesodesma donacium* (Bivalvia). *J. Shellfish Res.* 26, 43–49.
- Ries, J., Cohen, A., McCorkle, D., 2010. A nonlinear calcification response to CO₂-induced ocean acidification by the coral *Oculina arbuscula*. *Coral Reefs* 29, 661–674.
- Smith, S.V., 1973. Carbon dioxide dynamics: a record of organic carbon production, respiration, and calcification in the Eniwetok reef flat community. *Limnol. Oceanogr.* 18, 106–120.
- Stromgren, T., 1976. Skeleton growth of the hydrocoral *Millepora complanata* Lamarck in relation to light. *Limnol. Oceanogr.* 21, 156–160.
- Sun, T.C., Mori, S., Roper, J., Brown, C., Hooser, T., Burr, D.B., 1992. Do different fluorochrome labels give equivalent histomorphometric information? *Bone* 13, 443–446.
- Tambutte, E., Allemand, D., Bourge, I., Gattuso, J.P., Jaubert, J., 1995. An improved ⁴⁵Ca protocol for investigating physiological mechanisms in coral calcification. *Mar. Biol.* 122, 453–459.
- Tambutte, E., Tambutte, S., Segonds, N., Zoccola, D., Venn, A., Erez, J., Allemand, D., 2012. Calcein labelling and electrophysiology: insights on coral tissue permeability and calcification. *Proc. R. Soc. B: Biol. Sci.* 279, 19–27.
- Thebault, J., Chauvaud, L., Clavier, J., Fichez, R., Morize, E., 2006. Evidence of a 2-day periodicity of striae formation in the tropical scallop *Comptopallium radula* using calcein marking. *Mar. Biol.* 149, 257–267.
- Thorrold, S., Jones, G.P., Hellberg, M.E., Burton, R.S., Swearer, S.E., Neigel, J., Morgan, S.G., Warner, R.R., 2002. Quantifying larval retention and connectivity in marine populations with artificial and natural markers. *Bull. Mar. Sci.* 70s, 291–308.
- Thorrold, S.R., Jones, G.P., Planes, S., Hare, J.A., 2006. Transgenerational marking of embryonic otoliths in marine fishes using barium stable isotopes. *Can. J. Fish. Aquat. Sci.* 63, 1193–1197.
- Tribollet, A., Godinot, C., Atkinson, M., Langdon, C., 2009. Effects of elevated pCO₂ on dissolution of coral carbonates by microbial euendoliths. *Global Biogeochem. Cycles* 23, GB3008.
- Trotter, J., Montagna, P., McCulloch, M., Silenzi, S., Reynaud, S., Mortimer, G., Martin, S., Ferrier-Pages, C., Gattuso, J.-P., Rodolfo-Metalpa, R., 2011. Quantifying the pH vital effect in the temperate zooxanthellate coral *Cladocora caespitosa*: validation of the boron seawater pH proxy. *Earth Planet. Sci. Lett.* 303, 163–173.
- Vago, R., Gill, E., Collingwood, J.C., 1997. Laser measurements of coral growth. *Nature* 386, 30–31.
- Venn, A.A., Tambutté, E., Holcomb, M., Allemand, D., Tambutté, S., 2011. Live tissue imaging shows reef corals elevate pH under their calcifying tissue relative to seawater. *PLoS One* 6 (5), e20013. <http://dx.doi.org/10.1371/journal.pone.0020013>.
- Wilson, C.A., Beckman, D.W., Dean, J.M., 1987. Calcein as a fluorescent marker of otoliths of larval and juvenile fish. *Trans. Am. Fish. Soc.* 116, 668–670.
- Zar, J.H., 1984. *Biostatistical Analysis*. Prentice Hall, Englewood Cliffs, NJ.