




Contents lists available at ScienceDirect

Harmful Algae

journal homepage: www.elsevier.com/locate/hal



Modified local sands for the mitigation of harmful algal blooms

Gang Pan^{a,*}, Jing Chen^b, Donald M. Anderson^c

^a Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences, Beijing 10085, China

^b Yantai Institute of Coastal Zone Research, Chinese Academy of Sciences, Yantai 264003, China

^c Biology Department, Woods Hole Oceanographic Institution, MS 32, Woods Hole, MA 02543, USA

ARTICLE INFO

Article history:

Received 8 July 2010

Received in revised form 27 January 2011

Accepted 27 January 2011

Available online 2 February 2011

Keywords:

Chitosan

Harmful algal bloom

Modified sands

Polyaluminum chloride (PAC)

Seawater

Synergistic effect

ABSTRACT

A new method was developed for marine harmful algal bloom (HAB) mitigation using local beach sand or silica sand modified with chitosan and polyaluminum chloride (PAC). Untreated sand was ineffective in flocculating algal cells, but 80% removal efficiency was achieved for *Amphidinium carterae* Hulburt and *Chlorella* sp. in 3 min ($t_{80} = 3$ min) using 120 mg L⁻¹ sand modified with 10 mg L⁻¹ PAC and 10 mg L⁻¹ chitosan. After several hours 92–96% removal was achieved. The t_{80} for removing *A. carterae* using the modifiers only (PAC and chitosan combined) was 60 min and for *Chlorella* sp. 120 min, which are much slower than with the corresponding modified sand. Sands were critical for speeding up the kinetic processes of flocculation and sedimentation of algal flocs. PAC was helpful in forming small flocs and chitosan is essential to bridge the small flocs into large dense flocs. Chitosan was also important in inhibiting the escape of cells from the flocs. Chitosan and PAC used together as modifiers make it possible to use local beach sands for HAB mitigation in seawater. Economical and environmental concerns could be reduced through the use of sands and biodegradable chitosan, but the potential impacts of PAC need further study.

© 2011 Elsevier B.V. All rights reserved.

1. Introduction

Harmful algal blooms (HABs) pose a serious threat to public health, aquatic organisms, commercial fisheries, and the quality of freshwater lakes, rivers and reservoirs, as well as marine coastal environments. Over the past decade, there has been increasing interest in bloom mitigation strategies, though progress towards field applications has still been slow (Anderson, 1997). Significant attention has been focused on the use of clays as a means to remove HAB cells from the water column through flocculation and sedimentation. Many of these experiments were laboratory based (Beaulieu et al., 2005; Pan et al., 2006a; Pierce et al., 2004; Sengco et al., 2001; Yu et al., 1994), with some field demonstrations in Japan (Shirona, 1989), Australia (Atkins et al., 2001), China (Pan et al., 2006b) and South Korea (e.g., Lee et al., 2008). The environmental impacts of clay flocculation are generally positive, though there are studies that document negative effects. On the positive side, clay flocculation had little or no effect on marine organisms such as juvenile clams, fish, and invertebrates (Lewis et al., 2003; Archambault et al., 2004; Sengco and Anderson, 2004).

In one of these studies, however, a growth effect on juvenile hard clams was observed (compared to no-clay controls) with clay maintained in suspension for two weeks. These results suggest that clay applications in the field are likely more detrimental to clams under flow conditions leading to prolonged in situ resuspension of clay than under conditions that promote rapid sedimentation. Shumway et al. (2003) also report negative impacts on filter-feeding invertebrates using relatively high levels of clay. The magnitude of impacts is thus dependent on the flow regime, duration of exposure to resuspended clay, and the total clay loading.

However, clays are not immediately available at some locations that have HAB problems, and transportation costs may render this method uneconomical. There is also a common ecological concern about the dumping of large amounts of exotic materials into aquatic systems. As an alternative strategy, the use of native ecological materials such as local beach sands or soil (that naturally enter the aquatic system through rivers or rainfall) could in principle minimize the costs and ecological risk to aquatic environments. Sands, however, have markedly different physical characteristics from clays, and by themselves, will not flocculate and remove HAB cells.

In freshwater HAB mitigation, Pan and co-workers found that local soil particles including sands can be highly effective in removing cyanobacterial cells and improving water quality, but

* Corresponding author. Tel.: +86 10 62849686; fax: +86 10 62923541.

E-mail addresses: gpan@rcees.ac.cn, gpan6@yahoo.co.uk (G. Pan), jingchen2004126@126.com (J. Chen), danderson@whoi.edu (D.M. Anderson).

only after modification using small amounts of a natural, biodegradable material called chitosan (Pan et al., 2006b; Zou et al., 2006; Pan et al., 2011). These authors found that the polymeric netting and bridging function of chitosan was the key mechanism that allowed local soil particles to be highly effective in flocculating HAB cells. In this approach, the chitosan made a “net” that captured the HAB cells and other particles, and the soils provided the ballast or mass to carry the aggregates to the bottom. These encouraging results in freshwater have, however, limited direct applicability in marine systems, as high ionic strength and alkalinity prevent the unfolding of the polymer chain, thereby weakening chitosan's netting and bridging properties (Qun and Ajun, 2006; Zou et al., 2005).

Polyaluminum chloride (PAC), a commonly used inorganic coagulant, is highly effective in potable water treatment where it is used routinely to flocculate and remove suspended particles. PAC has been tested in marine systems and has been shown to reduce the amount of clays needed to remove HAB organisms (Pierce et al., 2004; Sengco et al., 2001; Yu et al., 1994). The addition of PAC increases the chemical affinity of clay surfaces. According to laboratory studies, however, algal cell flocculation by clays plus PAC was temporary (Sengco et al., 2001; Sun et al., 2004). Most of the cells could escape from the flocs and resume their growth. Motile dinoflagellate species were thus more difficult to be removed permanently through flocculation compared to non-motile diatoms (Yu et al., 1994), indicating that motility was an important factor affecting bloom mitigation through clay flocculation. Furthermore, the PAC floc was light, which did not settle easily or was resuspended with only modest currents (Beaulieu et al., 2005).

No efforts have been made thus far to use local beach sands to irreversibly flocculate and sediment marine HAB cells. Here, a modification of the approach to suppress freshwater HABs using local beach sands and polymers was developed for algal bloom mitigation in seawater. The synergistic effects of chitosan and PAC (hereafter termed “modifiers”) with two types of sands were investigated for the removal of *Amphidinium carterae* and *Chlorella* sp. The results demonstrate that it is possible to use modified local or commercially available sands to irreversibly remove a high percentage of the two types of HAB cells from seawater.

2. Materials and methods

2.1. Algal species and culture

Two algal species were used – *A. carterae* Hulburt, a motile dinoflagellate, and a marine *Chlorella* sp. which is very small, and non-motile. *A. carterae* is considered a HAB species because of its production of haemolysins, and it has also been linked to fish mortalities (Hulburt, 1957; Yasumoto et al., 1987). Although *Chlorella* is not listed as a harmful species on some lists, it is known for its ability to produce dense blooms that can have adverse consequences, such as the decimation of the oyster industry on Long Island following eutrophication stimulated by duck farm effluents (Ryther, 1954). *A. carterae* was obtained from Oceanography College, Ocean University of China and *Chlorella* sp. was supplied by Seaweed Inheritance Breeding Center of Shandong Oriental Ocean Sci.-Tech. Co. Ltd.

The cells were grown in f/2 medium (Guillard and Hargraves, 1993) made with synthetic seawater. The synthetic seawater was composed of 23.939 g L⁻¹ NaCl, 5.079 g L⁻¹ MgCl₂·6H₂O, 3.994 g L⁻¹ Na₂SO₄, 1.123 g L⁻¹ CaCl₂, 0.667 g L⁻¹ KCl, 0.196 g L⁻¹ NaHCO₃, 0.098 g L⁻¹ KBr, 0.027 g L⁻¹ H₃BO₃, 0.003 g L⁻¹ NaF and 0.024 g L⁻¹ SrCl₂·6H₂O. The medium was adjusted to pH 8.2 before autoclaving by adding either 0.1 mol L⁻¹ NaOH or 0.1 mol L⁻¹ HCl solutions.

Algal batch cultures were maintained at 25 ± 1 °C under continuous cool white fluorescent light of 2000–3000 lux on a 12 h light and 12 h darkness regimen in the illuminating incubator (LRH-250-G, Guangdong Medical Apparatus Co., Ltd., China).

2.2. Sands and modifiers

Two kinds of sand were used. One was SiO₂ (silica sand) analytical grade, purchased from Sinopharm Chemical reagent Co., Ltd. Another was local sand collected from a Yellow Sea beach in Yantai, China. The two sands were washed with deionized water, dried at 100 °C, and sieved through 180 mesh (<90 µm).

Chitosan was obtained from Qingdao Haisheng Bioengineering Co., Ltd. The chitosan flakes were dissolved by adding 100 mg chitosan to 10 mL of 0.5% HAc and stirring until all the chitosan was dissolved. This solution was diluted with deionized water to obtain a final concentration of 1 mg mL⁻¹ before use (Zou et al., 2006). PAC was supplied by Dagang Reagent Plant, Tianjin, China. The basicity (B = [OH]/[Al]) of PAC was 2.4 and its Al₂O₃ content was 30%. The PAC was dissolved in deionized water to obtain a solution of 1 mg mL⁻¹. The chitosan and PAC solutions were prepared freshly before each set of experiments.

2.3. Algal flocculation

Flocculation experiments were conducted in a jar test apparatus (ZR3-6, Zhongrun Water Industry Technology Development Co., Ltd., China) using cultures in mid- to late-exponential growth phase. The initial cell concentrations of *A. carterae* and *Chlorella* sp. were 3.25–3.42 × 10⁵ cells mL⁻¹ and 6.65–6.82 × 10⁶ cells mL⁻¹, respectively. Two hundred milliliters of experimental culture were transferred into a 250 mL beaker, stirred at 200 rpm for 2 min, followed by 30 rpm for another 5 min. Chitosan alone, PAC alone, chitosan plus PAC together, and chitosan plus PAC plus sands were added to the algal culture in different flocculation experiments. The control culture was run without adding any sands or modifiers.

Samples from 2 cm below the surface of the experimental beaker were collected after sedimentation at different times and the cells enumerated in a counting chamber under an electromotive microscope (Axioskop 2 mot plus, Carl ZEISS, Germany) after being fixed by Lugol solution. The removal efficiency of cells was calculated as (initial cell concentration – sample cell concentration)/initial cell concentration × 100%. Algal flocs were collected by pipette and observed under the microscope.

Algal floc size and size distribution during the flocculation process were monitored with a laser particle size analyzer Mastersizer 2000 (Malvern Co., United Kingdom). The culture was drawn into the Mastersizer and back to the jar by a peristaltic pump (BT00-300M, Baoding Longer Precision Pump Co., Ltd., China) at a flow rate of 34 mL min⁻¹ (Zhang et al., 2007). Samples were at the same position in the jar, which was located between the impeller and the top of suspension. Algal floc size was denoted by the measured mean diameter (d_{50}).

2.4. Viability and growth of algae after flocculation

The effect of PAC or chitosan with PAC on the viability and the growth of *A. carterae* after flocculation was investigated using two strategies. In the first experiment, fresh f/2 medium was added to the supernatant without disturbing the algal flocs (Sengco et al., 2001; Sun and Choi, 2004). This flask was maintained in an illuminated incubator, and viability and growth of the cells were monitored by measuring the cell concentrations in the supernatant after 24 and 48 h. In the second experiment, flocs were maintained in the incubator without fresh f/2 medium or light.




Fig. 1. Algal removal efficiency of 100 mg L⁻¹ local sands, 100 mg L⁻¹ silica sands, modifiers (10 mg L⁻¹ chitosan plus 10 mg L⁻¹ PAC), modified local sands (10 mg L⁻¹ chitosan plus 10 mg L⁻¹ PAC plus 100 mg L⁻¹ local sands) and modified silica sands (10 mg L⁻¹ chitosan plus 10 mg L⁻¹ PAC plus 100 mg L⁻¹ silica sands) at different time. (A) *A. carterae* and (B) *Chlorella* sp.

3. Results

3.1. Algal flocculation using modified sands

Compared with control experiments, 100 mg L⁻¹ silica sand or local sand was ineffective in removing *A. carterae* and *Chlorella* sp. (Fig. 1). However, sands modified using chitosan and PAC combined were highly efficient in flocculating and sinking algal cells. The removal efficiency with 120 mg L⁻¹ modified sands containing 10 mg L⁻¹ chitosan and 10 mg L⁻¹ PAC reached 80% for the two algal species within 3 min ($t_{80} = 3$ min), whereas the removal efficiencies of only 10 mg L⁻¹ chitosan plus 10 mg L⁻¹ PAC on *A. carterae* (Fig. 1A) and *Chlorella* sp. (Fig. 1B) were 54% and 43%, respectively. The t_{80} of the modifiers alone for *A. carterae* removal was 60 min and that for *Chlorella* sp. was 120 min. Using only sands, the removal efficiencies of *A. carterae* and *Chlorella* sp. after 240 min were 26% and 7% (Fig. 1A and B). This increased to 96% and 92% when the chitosan and PAC modifiers were added with the sand. The results in Fig. 1 also demonstrate that there was no large difference between silica sand and local beach sand on HAB cell removal if the modifiers chitosan and PAC were present.

3.2. Synergistic effect of chitosan and PAC on algal cell removal

When chitosan was used alone, cell removal efficiencies increased with increasing dosage of chitosan (0–20 mg L⁻¹ for *A. carterae* and 0–50 mg L⁻¹ for *Chlorella* sp.; Fig. 2). However, the removal efficiency of *A. carterae* (Fig. 2A) was maximally 71% at

20 mg L⁻¹ chitosan and that of *Chlorella* sp. (Fig. 2B) was only 51% at 50 mg L⁻¹, which suggests that chitosan is not as efficient at removing algal cells from seawater as it is in fresh water (Pan et al., 2006b; Zou et al., 2006).

Cell removal efficiency for both species increased when PAC and chitosan were used together (Fig. 2). After the addition of 5 mg L⁻¹ PAC with 10 mg L⁻¹ chitosan, the removal efficiency of *A. carterae* and *Chlorella* sp. increased to 92% and 62% from 68% and 11%, respectively. When 10 mg L⁻¹ PAC was added with 10 mg L⁻¹ chitosan, the *A. carterae* removal efficiency increased by an additional 28% over that with chitosan alone, and that of *Chlorella* sp. increased by 78%.

3.3. Synergistic effect of chitosan and PAC on algal floc formation

The formation and development of algal flocs using 10 mg L⁻¹ PAC or PAC with 10 mg L⁻¹ chitosan were investigated using *Chlorella* sp. as the target species. The floc size (Fig. 3A) and size distributions (Fig. 3B) were monitored. Compared with PAC alone, the algal flocs of PAC plus chitosan increased in size much faster in the first 2 min. During the slow stir phase, algal floc size increased to a plateau. The floc size of PAC plus chitosan increased to 860 μ m, compared to that of PAC alone, for which the size was approximately 600 μ m. The floc produced by chitosan and PAC appeared rapidly and quickly increased in size to form larger particles than with PAC only.

At 7 min, the stir was over and floc size distribution curves were shown in Fig. 3B. The floc size distribution of PAC alone ranged




Fig. 2. Synergistic effect of chitosan and PAC on algae removal. (A) *A. carterae* and (B) *Chlorella* sp.




Fig. 3. Synergistic effect of chitosan and PAC on algal flocs. (A) Floc size and (B) floc size distributions at 7 min.

between 316 μm and 1259 μm, with the highest peak at 631 μm. The size distribution of PAC plus chitosan was between 417 μm and 2188 μm, with the highest peak at 955 μm.

3.4. Synergistic effect of chitosan and PAC on cell viability

An experiment examining the synergistic effect of chitosan and PAC on the viability and growth of *A. carterae* was divided into three treatments: (1) 10 mg L⁻¹ PAC only, (2) 10 mg L⁻¹ PAC plus 10 mg L⁻¹ chitosan, and (3) 10 mg L⁻¹ PAC plus 20 mg L⁻¹ chitosan. After these flocculation experiments, the residual cell concentration in the supernatant of the three treatments was 1.2–1.6 × 10⁴ cells mL⁻¹, approximately 4% of the original concentration prior to the treatment. The cell concentration for all the treatments roughly doubled to 2.8–3.0 × 10⁴ cells mL⁻¹ after 24 h of incubation in an incubator with light and added nutrients (Fig. 4A). After another 24 h, the cell concentration with PAC only increased dramatically to 12.4 × 10⁴ cells mL⁻¹, while the concentration in the treatments of PAC plus 20 mg L⁻¹ chitosan rose to 5.05 × 10⁴ cells mL⁻¹, approximately half of the concentration with PAC only.

The results shown in Fig. 4B demonstrate that the cell concentration in the supernatant of the three treatments in the incubator with no light or added nutrients decreased gradually throughout the study interval. However, the algal cell concentrations of PAC plus chitosan used together were less than that of PAC alone and the cell concentration was inversely related to the chitosan dosage. After 28 days, the concentration of algal cells in supernatant was only 300 cells mL⁻¹, indicative of almost no recovery of *A. carterae* cells under conditions similar to those found near bottom sediments.




Fig. 4. Synergistic effect of chitosan and PAC on algae viability. (A) With light and added nutrients and (B) with no light or added nutrients.

4. Discussion

In this study, a method was developed that uses sands or local soils that could be collected from the immediate vicinity of a HAB, and used in conjunction with small amount of chitosan and PAC to flocculate and effectively remove cells from the water column. Our results demonstrate that PAC was needed to maintain the netting and bridging function of chitosan in seawater and to form small flocs, while chitosan was essential in bridging the small flocs into large and dense flocs that hindered the escape of cells from the flocs. As the safe and cheap carrier of these modifiers, sand was critical for speeding up sedimentation. This approach, which was a modification of the one used successfully for HAB removal in freshwater systems (Pan et al., 2006b; Pan et al., 2011), greatly minimizes environmental concerns for mitigation of HABs in seawater using clays since the use of native beach sands has few environmental concerns. As discussed below, however, there are still some issues that need to be addressed if this method is used for field applications on natural blooms.

4.1. Synergistic effects of chitosan plus PAC

The flocculation of algal cells in natural waters occurs as a result of attractive anion–cation interactions, as well as hydrophobic or polymer interactions (Divakaran and Pillai, 2001; Strand et al., 2002). Sands alone are much less efficient in flocculating algal cells compared to clays such as kaolinite, montmorillonite, and sepiolite (Pan et al., 2006a,b; Pierce et al., 2004; Sengco et al., 2001; Yu et al., 1994). Chitosan and PAC as modifiers increase the surface charge of sands and enhance the netting and bridging interactions with algal

cells. Sands also provide the mass or ballast to carry flocs to bottom sediments.

Chitosan, a cellulose-like polyelectrolyte biopolymer, is derived from the alkaline deacetylation of crustacean chitin, which possesses several intrinsic characteristics of coagulants and flocculants, i.e., high cationic charge density, long polymer chains, bridging of aggregates and precipitation (Renault et al., 2009; Rinaudo, 2006). Chitosan, by itself, does not flocculate effectively in seawater (Fig. 2). This is because its molecular structure includes abundant amino groups ($-\text{NH}_2$) and hydroxyl groups ($-\text{OH}$) on the chain. The active amine group ($-\text{NH}_2$) of chitosan is easily protonated as $-\text{NH}_3^+$ in dilute acidic solutions, and there is a strong electrostatic repulsion force within and between molecules (Rinaudo, 2006). The high content of positively charged amine groups in the chitosan structure facilitates electrostatic interactions between polymer chains and negatively charged contaminants (Huang et al., 2000; Renault et al., 2009). However, in high ionic strength solutions such as seawater, counter-ions accumulate near the $-\text{NH}_3^+$ group, which would screen the protonated amine groups and decrease the electrostatic repulsion among them (Qun and Ajun, 2006; Schatz et al., 2003). This prevents the unfolding of the molecular chain, thereby weakening its netting and bridging properties (Zou et al., 2005).

In contrast to chitosan, the high ionic strength of seawater is beneficial to PAC flocculation due to the reduction of the thickness of the electrical double layer which enhances the collision probability of granules. PAC supplies cationic hydrolysis products that are strongly adsorbed on negative particles and can give effective destabilization, leading to the formation of micro-flocs (Renault et al., 2009). Particles with thinner electrical double layers are easier to coagulate because of reduced repulsion. With the high salinity of seawater, flocculation of particles is increased because the thickness of the electrical double layer is decreased due to the compression of the electrolytes (Han and Kim, 2001; Pan et al., 2006b). This explains why PAC is effective in flocculating HAB cells in seawater and why the algal cell removal efficiencies of chitosan are increased remarkably with the addition of PAC. PAC cannot be used by itself in seawater, however, since, discussed by Beaulieu et al. (2005), PAC flocs are light and fluffy and do not settle even in light flow regimes. If these small flocs can be combined and form a stronger, larger, and heavier flocs, then the limitations of PAC flocs can be overcome.

The amino groups ($-\text{NH}_2$) and hydroxyl groups ($-\text{OH}$) in chitosan's molecular structure contain single-pair electrons that can offer the electron pair to empty trajectories of metal ions; they then chelate into a complex compound (Bassi et al., 2000). It was reported that there was a positive correlation between chitosan

and PAC and the effect of chitosan adsorbing Al^{3+} in solution was very obvious (Zeng et al., 2008). The cationic hydrolysis products of PAC that are adsorbed on the molecule chain of chitosan might increase electrostatic repulsion between them and protonated groups ($-\text{NH}_3^+$), which would in turn be beneficial to the unfolding of chitosan's molecular chain and weaken the negative effect of high ionic strength on chitosan's netting and bridging properties in seawater. Therefore, PAC and chitosan are complementary in flocculating HAB cells in seawater. Larger and denser algal flocs are formed by the compression of electrical double layer, charge neutralization, adsorption, and netting interactions to bind and bridge cells tightly.

4.2. Cell escape from flocs

As shown in Fig. 4, with light and nutrients provided to cells flocculated using PAC and chitosan alone, cell concentrations in the supernatant doubled in 24 h, and then doubled again 24 h later. *Amphidinium* can grow rapidly, with growth rates as high as 2.7 divisions per day (Ismael et al., 1999), so the cell increase in the supernatant of the chitosan plus PAC treatment could be explained entirely by growth with little or no contribution from cells escaping from the flocs. The much larger increase in cell abundance in the PAC only treatment suggests that a significant number of cells escaped into the supernatant.

Chitosan flocs were fibrous and formed large entangled masses resembling cobwebs by bridging mechanisms (Fig. 5A). The protonated amine group of chitosan attract negatively charged algal cells to produce large and complex flocs that help to prevent the escape of motile cells. In contrast, the flocs of PAC alone were small and there were large numbers of cells around the flocs (Fig. 5B). This implies that PAC does not bridge the algal cells firmly nor bind them as strongly as chitosan does. Overall, the number of cells escaping from the PAC plus chitosan flocs was small, and the method appeared promising for bloom mitigation. The addition of sand would make cell escape even more difficult.

4.3. Environmental impacts

One of the challenging and controversial aspects of HAB research relates to methods to directly control or suppress blooms (Anderson, 1997). Of the many methods that have been proposed, removal of HAB cells through clay flocculation is seen by some as promising in terms of efficiency, cost, and environmental impacts (e.g., Sengco and Anderson, 2004; Lee et al., 2008). There are, however, those who feel that the environmental impacts of this approach are unacceptable, or poorly understood. In addition to




Fig. 5. Algal flocs micrographs with the magnification of 50 $\times$. (A) Chitosan and *A. carterae* and (B) PAC and *A. carterae*.

the possible adverse ecological impact caused by the addition of large amount of exotic materials (Shumway et al., 2003), other concerns expressed relates to the constituents in the clay, which might include nutrients such as phosphorus, or toxic or harmful metals and radioactive materials bound to the clay. As an alternative to clays, sands are relatively inert or refractory and thus may minimize these impacts. Most importantly, as a native part of the ecosystem, beach sand is ecologically safe to the marine system which may avoid the fundamental concern associated with clays. Large-scale dredging and beach nourishment projects abound in nearshore waters worldwide, suggesting that environmental opposition to HAB mitigation efforts using local sands might be minimal. In cases where beach sands need to be conserved, commercially available sands may also be safe, cheap and easily available to be used.

The modification technique using chitosan and PAC can not only turn local beach sands or local soils into highly effective flocculants in the mitigation of HABs in seawater, but is also useful in reducing the loading of sands/soils required for effective cell removal, which is crucial for large scale field applications. Chitosan, a commercially available product of edible food additives, is known to be a biodegradable and non-toxic natural polymer. Compared with other chemical reagents, chitosan is environmental friendly, but it might be a source of oxygen demand as it decays. The amount of chitosan used is, however, much less than the amount of algal biomass being sedimented, so this is not a serious concern. Nevertheless, it may be worthwhile to develop techniques that could carry and release oxygen with the flocs to combat this potential problem (Pan et al., 2009). In some coastal areas, it is also possible to sink the algal blooms into the bottom and cover them using a second layer of sands or local soils so that the cells can be permanently buried and sealed in the sediment and turned into fertilizers for the growth of seaweeds, as Pan et al. (2011) demonstrated in shallow lakes. By decomposing the algal cells and the modifiers and converting them into the biomass of seaweeds, the harmful blooms may be turned into useful resources for the improvement of the ecosystem. However, this possibility needs further study in marine systems affected by HABs. Although PAC (a compound used in drinking water treatment) was needed to maintain the netting and bridging function of chitosan in seawater, the adverse ecological effects of this compound in seawater remain a concern. More research is needed in this area before larger-scale applications can be undertaken. Similarly, efforts are needed to identify new, environmentally benign modifiers that could replace PAC in this bloom control strategy.

5. Conclusion

Dispersal of sands or local soils modified with chitosan and PAC achieved high removal efficiency of marine HAB cells in a short time and prevented the escape of significant numbers of motile organisms from the algal flocs. This method greatly reduces potential environmental impacts by using relatively inert or refractory sand or local soils and by using a biodegradable polymer such as chitosan, but there may be environmental concerns about the use of PAC. With some additional studies, this approach shows great promise to become an effective and environmentally acceptable strategy for HAB mitigation.

Acknowledgements

The research was funded by the National Key Project for Basic Research (2008CB418105, 2010CB933600), for which the authors are grateful. Support for DMA was provided by GOMTOX program through NOAA Grant NA06NOS4780245. Additional support came from NSF grant OCE-0430724, DMS-0417769 and NIEHS grant

1P50-ES01274201 (Woods Hole Center for Oceans and Human Health). [SS]

References

- Anderson, D.M., 1997. Turning back the harmful red tide. *Nature* 388, 513–514.
- Archambault, M.C., Bricelj, V.M., Grant, J., Anderson, D.M., 2004. Effects on juvenile hard clams, *Mercenaria mercenaria*, from the application of phosphatic clays to mitigate harmful algal blooms. *Mar. Biol.* 144, 553–565.
- Atkins, R., Rose, T., Brown, R.S., Robb, M., 2001. The microcystis cyanobacteria bloom in the Swan River—February 2000. *Water Sci. Technol.* 43 (9), 107–114.
- Bassi, R., Prasher, S.O., Simpson, B.K., 2000. Removal of selected metal ions from aqueous solutions using chitosan flakes. *Sep. Sci. Technol.* 35 (4), 547–560.
- Beaulieu, S.E., Sengco, M.R., Anderson, D.M., 2005. Using clay to control harmful algal blooms: deposition and resuspension of clay/algal flocs. *Harmful Algae* 4 (1), 123–138.
- Divakaran, R., Pillai, V.N.S., 2001. Flocculation of kaolinite suspensions in water by chitosan. *Water Res.* 35 (16), 3904–3908.
- Guillard, R.R.L., Hargraves, P.E., 1993. *Stichochrysis-immobilis* is a diatom, not a chrysophyte. *Phycologia* 32 (3), 234–236.
- Han, M.Y., Kim, W., 2001. A theoretical consideration of algae removal with clays. *Microchem. J.* 68 (2–3), 157–161.
- Huang, C., Chen, S., Ruhsing Pan, J., 2000. Optimal condition for modification of chitosan: a biopolymer for coagulation of colloidal particles. *Water Res.* 34 (3), 1057–1062.
- Hulburt, E.M., 1957. The taxonomy of unarmored Dinophyceae of shallow embayments off Cape Cod, Massachusetts. *Biol. Bull.* 112 (2), 196–219.
- Ismael, A.A.H., Halim, Y., Khalil, A.G., 1999. Optimum growth conditions for *Amphidinium carterae* Hulburt from eutrophic waters in Alexandria (Egypt) and its toxicity to the brine shrimp *Artemia salina*. *Grana* 38 (2–3), 179–185.
- Lee, Y.-J., Choi, J.-K., Kim, E.-K., Youn, S.-H., Yang, E.-J., 2008. Field experiments on mitigation of harmful algal blooms using a Sophorolipid–Yellow clay mixture and effects on marine plankton. *Harmful Algae* 7 (2), 154–162.
- Lewis, M.A., Dantin, D.D., Walker, C.C., Kurtz, J.C., Greene, R.M., 2003. Toxicity of clay flocculation of the toxic dinoflagellate, *Karenia brevis* to estuarine invertebrates and fish. *Harmful Algae* 2 (4), 235–246.
- Pan, G., Yang, B., Li, L., Liao, L., Ding, C., Wang, D., 2009. A method for anaerobic sediments remediation and ecological restoration in eutrophic shallow lakes. Chinese Patent Application No.: 200910080563.5.
- Pan, G., Zhang, M., Chen, H., Zou, H., Yan, H., 2006a. Removal of cyanobacterial blooms in Taihu Lake using local soils. I. Equilibrium and kinetic screening on the flocculation of *Microcystis aeruginosa* using commercially available clays and minerals. *Environ. Pollut.* 141 (2), 195–200.
- Pan, G., Zou, H., Chen, H., Yuan, X., 2006b. Removal of harmful cyanobacterial blooms in Taihu Lake using local soils III. Factors affecting the removal efficiency and an in situ field experiment using chitosan-modified local soils. *Environ. Pollut.* 141 (2), 206–212.
- Pan, G., Yang, B., Wang, D., Chen, H., Tian, B.H., Zhang, M.L., Yuan, X.Z., Chen, J., 2011. In-lake algal bloom removal and submerged vegetation restoration using modified local soils. *Ecol. Eng.* 37, 302–308.
- Pierce, R.H., Henry, M.S., Higham, C.J., Blum, P., Sengco, M.R., Anderson, D.M., 2004. Removal of harmful algal cells (*Karenia brevis*) and toxins from seawater culture by clay flocculation. *Harmful Algae* 3 (2), 141–148.
- Qun, G., Ajun, W., 2006. Effects of molecular weight, degree of acetylation and ionic strength on surface tension of chitosan in dilute solution. *Carbohydr. Polym.* 64 (1), 29–36.
- Renault, F., Sancey, B., Badot, P.M., Crini, G., 2009. Chitosan for coagulation/flocculation processes—an eco-friendly approach. *Eur. Polym. J.* 45 (5), 1337–1348.
- Rinaudo, M., 2006. Chitin and chitosan: properties and applications. *Prog. Polym. Sci.* 31 (7), 603–632.
- Ryther, J.H., 1954. The ecology of phytoplankton blooms in Moriches Bay and Great South Bay, Long Island, New York. *Biol. Bull.* 106 (2), 198–209.
- Schatz, C., Viton, C., Delair, T., Pichot, C., Domard, A., 2003. Typical physicochemical behaviors of chitosan in aqueous solution. *Biomacromolecules* 4 (3), 641–648.
- Sengco, M.R., Anderson, D.M., 2004. Controlling harmful algal blooms through clay flocculation. *J. Eukaryot. Microbiol.* 51 (2), 169–172.
- Sengco, M.R., Li, A., Tugend, K., Kulis, D., Anderson, D.M., 2001. Removal of red- and brown-tide cells using clay flocculation. I. Laboratory culture experiments with *Gymnodinium breve* and *Aureococcus anophagefferens*. *Mar. Ecol. Prog. Ser.* 210, 41–53.
- Shirona, A., 1989. Red tide problem and countermeasures. *Int. J. Aquat. Fish. Technol.* 1, 195–223.
- Shumway, S.E., Frank, D.M., Ewart, L.M., EvanWard, J., 2003. *Aquaculture Res.* 34, 1391–1402.
- Strand, S.P., Nordengen, T., Ostgaard, K., 2002. Efficiency of chitosans applied for flocculation of different bacteria. *Water Res.* 36 (19), 4745–4752.
- Sun, X.X., Choi, J.K., 2004. Recovery and fate of three species of marine dinoflagellates after yellow clay flocculation. *Hydrobiologia* 519 (1–3), 153–165.
- Sun, X.X., Lee, Y.J., Choi, J.K., Kim, E.K., 2004. Synergistic effect of sophorolipid and loess combination in harmful algal blooms mitigation. *Mar. Pollut. Bull.* 48 (9–10), 863–872.
- Yasumoto, T., Seino, N., Murakami, Y., Murata, M., 1987. Toxins produced by benthic dinoflagellates. *Biol. Bull.* 172 (1), 128–131.
- Yu, Z.-M., Zou, J.-Z., Ma, X.-N., 1994. Application of clays to removal of red tide organisms: II. Coagulation of different species of red tide organisms with

- montmorillonite and effect of clay pretreatment. *Chin. J. Oceanol. Limnol.* 12 (4), 316–324.
- Zeng, D., Wu, J., Kennedy, J.F., 2008. Application of a chitosan flocculant to water treatment. *Carbohydr. Polym.* 71 (1), 135–139.
- Zhang, Z.G., Luan, Z.K., Zhao, Y., Cui, J.H., Chen, Z.Y., Li, Y.Z., 2007. Breakage and regrowth of flocs coagulation with polyaluminum chloride (PACl). *Environ. Sci.* 28 (2), 346–351.
- Zou, H., Pan, G., Chen, H., 2005. Effects of ionic strength on the flocculation and removal of cyanobacterial cells of *Microcystis aeruginosa* by clays. *Environ. Sci.* 26, 148–151.
- Zou, H., Pan, G., Chen, H., Yuan, X., 2006. Removal of cyanobacterial blooms in Taihu Lake using local soils II. Effective removal of *Microcystis aeruginosa* using local soils and sediments modified by chitosan. *Environ. Pollut.* 141 (2), 201–205.