

## Beaufort Gyre Exploration Project: Dispatch 8: Searching for the Perfect Ice

On August 20th 2003, at 7:30 AM *Louis S. St-Laurent's* position was 80° 00N, 150°00 W . This is our farthest North station and the turning point of our expedition.

During next two weeks, we will gradually move in the direction of Kugluktuk. There is still a lot of work waiting for us. The event of reaching the expedition's turning point also became a turning point in the weather and sea ice conditions. Pleasant, cooperative Arctic weather that had accompanied our work during the first two weeks, changed rapidly in the morning. We planned to deploy one of our four expendable buoys on August 21st, but wanted to make sure that the buoy would be installed on an old ice field with a thickness of about 2 meters so that it would work for at least one year, sending oceanic water temperature and salinity data to the Woods Hole Oceanographic Institution two times per day. Therefore, Scott Payment, the ship's Ice Observer, Adrian Godin, helicopter pilot, and WHOI scientist Andrey Proshutinsky executed a 90-minute ice reconnaissance helicopter flight to search for an appropriate ice field.

No good ice fields were found, and fog, clouds and a general decrease of visibility forced us to return to the ship. We decided that the search would be repeated during the 1-2 days, hoping that winds and currents would move the old and thick ice to the southeast – region of our future work.

*Last updated: October 19, 2015*



[Enlarge Image](#)

Helicopter pilot Adrian Godin inspects the helicopter before the sea ice reconnaissance flight.



[Enlarge Image](#)

The Ice Observer, Scott Payment is ready for the flight.



[Enlarge Image](#)

Andrey Proshutinsky (WHOI) gets instructions from the helicopter engineer, Steve Lloyd before take off.



[Enlarge Image](#)

Adrian Godin starts the helicopter's engine.



[Enlarge Image](#)

*Louis S. St-Laurent.*



[Enlarge Image](#)

Sea ice fields from an altitude of 400 meters. Some of the ice fields are huge (up to 10 miles in diameter, not shown here) but this is first-year ice with thickness less than 100 cm. In September and October, strong storms can destroy these fields and our buoy could sink. Stronger, thicker ice is needed.